8509

Melonax 7.5
Meloxicam Tablets 7.5 mg

Melonax 15
Meloxicam Tablets 15 mg

PRESENTATION
7.5mg/15mg: 10 tablets in a blister. 1x10 Tablets / 2x10 Tablets in blister in a carton.

COMPOSITION
Each tablet contains; Meloxicam BP 7.5mg/15mg
(**Common Excipients**) Microcrystalline Cellulose, Lactose, Croscarmellose Sodium, Maize Starch, Sodium Lauryl Sulphate, Magnesium Stearate.

PHARMACOLOGICAL ACTION
Melonax is a new Non-steroidal Anti-inflammatory drug (NSAID) of the oxicam class. It differs from other NSAIDs in that it has a greater selective inhibitory activity against the inducible isoform of cyclo oxygenase (COX-2) than against the COX-1 which is associated with gastrointestinal/renal adverse effects and inhibition of platelet aggregation effects of the NSAIDs.

INDICATIONS
Melonax is indicated in Osteoarthritis, Rheumatoid arthritis, low back pain, Dysmenorrhoea and other gynaecological disorder, dental and post-operative pain, in reducing pain of ear, nose and throat inflammation, pain of malignancy, sports injuries.

CONTRAINDICATION
Melonax is contraindicated in patients with known hypersensitivity. Active peptic ulcer, moderate to severe hepatic impairment, hypersensitivity to aspirin or other NSAIDs. It should not be given to patients who have experienced asthma, bronchospasm, urticaria or allergic type reactions after taking aspirin or other NSAIDs. Allergic and anaphylactoid reactions have been reported in patient receiving Meloxicam in clinical trials.

PRECAUTIONS
In/for patients with signs and/or symptoms suggesting liver dysfunction, or in whom an abnormal liver test has occurred should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with Meloxicam.
Caution should be used when initiating treatment with Meloxicam in patients with considerable dehydration.
Anaemia is sometimes seen in patients receiving Meloxicam. This may be due to fluid retention, GI blood loss.

ADVERSE REACTIONS
The most commonly reported reaction is gastrointestinal disturbances which included abdominal pain, diarrhea, dyspepsia, flatulence, nausea/vomiting. Less common gastrointestinal disturbances include colitis, peptic ulcer diseases, esophagitis, gastritis, GI bleeding.

ADVERSE EFFECTS
Nausea, vomiting, diarrhea, heart burn, epigastric pain, pruritus, skin rash, headache, somnolence and dizziness.

INTERACTIONS
Fenofibrate, salicyclic acid, valproic acid and *tolbutamide*: These drugs displace Meloxicam from the plasma binding sites.
Methotrexate and Frusemide: These drugs may be displaced from the plasma protein by Meloxicam
Warfarin: Exercise caution since efficacy may be increased by Meloxicam.
Theophylline: Efficacy of slow release theophylline preparations reduced by Meloxicam

DOSAGE AND ADMINISTRATION
As directed by the physician
Adults: Melonax tablets of 7.5mg one to two tablets are taken once daily for 7 days.
Melonax tablets of 15mg one tablet is taken once daily for 7 days.
It is administered through oral dosage.

STORAGE INSTRUCTIONS
Store below 30°C. Protect from light. Keep all medicines away from reach of children.

1040070

A product of **Strides Pharma Science Limited**
Strides House, Bilekahalli, Bannerghatta Road,
Bangalore - 560 076, Karnataka, India.

8509

------ 100 x 190 mm ------

Front Page Printing

Melonax 7.5
Meloxicam Comprimés 7.5 mg

Melonax 15
Meloxicam Comprimés 15 mg

PRÉSENTATION
7.5mg/15mg : Boite de 10 comprimés. 1 plaquette / 2 plaquettes de 10 comprimés dans une boite.

COMPOSITION
Chaque comprimé contient ; Meloxicam BP 7.5mg/15mg
(Excipients communs) cellulose microcristalline, lactose, sodium de Croscarmellose, fécule de maïs, sulfate laurique de sodium, stéarate de magnésium.

ACTION PHARMACOLOGIQUE
Melonax est un nouveau médicament anti-inflammatoire Non stéroïdiens (AINS) de la classe des oxicam. Il diffère des autres AINS parce qu'il a une plus grande activité inhibitrice sélective de la cycloxygénase (COX-2) que de la cycloxygénase COX-1 qui est associé aux effets nuisibles de type gastro-intestinale/rénale qui affecte l'action inhibitrice des AINS.

INDICATIONS
Melonax est indiqué dans les douleurs pelviennes.Ostéo-arthrites – arthrites rhumatoïdes – douleurs de dos – dysménorrhées et douleurs gynécologiques – douleurs dentaires et post chirurgicale – réduction des douleurs des oreilles
l'inflammation du nez et de la gorge – blessures sportives..

CONTRE-INDICATION
Melonax est contre-indique dans : Ulcère peptique actif, hypersensibilité à l'aspirine et autres AINS. Melonax est contre-indiqué aux patients qui ont déjà fait une crise d'asthme – un spasme bronchique – qui ont présenté un urticaire et une allergie à la prise de l'aspirine ou des autres AINS – des allergies et des réactions anaphylactiques ont été constaté chez des patients ayant pris le Melonax

PRÉCAUTIONS
Des patients ayant des signes ou des symptômes ayant trait avec un dysfonctionnement du foie doivent être examinés pour éviter des réactions hépatiques sévères pendant le traitement avec le Meloxicam. Des précautions doivent être prises avant d'initier un traitement à base de Meloxicam chez des patients déshydratés.Des patients recevant le Meloxicam ont souvent été anémiés. Cela pourrait être dû à la rétention des fluides et à la perte des globules du sang.L'anémie est parfois vue dans les patients recevant le Meloxicam. Ceci peut être dû à la conservation liquide, perte de globule de sang.

EFFETS SECONDAIRES
Les réactions indésirables les plus connues sont les troubles gastro – intestinaux qui comportent des douleurs abdominales, diarrhées, la dyspepsie, la flatulence, les nausées et les vomissements. Les troubles gastro-intestinaux moins importants qui peuvent survenir sont les colites, l'ulcère peptique, l'oesophagite, gastrite et les saignements.

EFFETS NUISIBLES
Nausée, vomissement, diarrhée, brûlures d'estomac, douleur épigastrique, prurit, éruption de peau, mal de tête, somnolence et vertige.

INTERACTIONS MEDICAMENTEUSES :
Le Fénofibrate, l'acide salicylique, l'acide valpolique, le Tolbutamide sont des molécules qui empêchent le Meloxicam de se fixer sur les sites plasmatiques.
Le Methotrexate et le Frusemide peuvent être
détachés de leurs sites plasmatiques par l'effet du Meloxicam.
Le Warfrarin exerce une précaution puisque son efficacité peut être accrue par le Meloxicam.

POSOLOGIE
D'Après la prescription du médecin
Adultes : Les comprimés de Melonax 15 mg se prennent à dose unique : Un comprimé par jour pendant 7 jours.
Les comprimés de Melonax 7.5mg se prennent : Un à deux comprimé par jour pendant 7 jours.
Il est administré par voie orale.

INSTRUCTIONS DE CONSERVATIONS : Le conserver en dessous de 30° C.
Le protéger contre la lumière. Le garder hors de la portée des enfants.

Un produit de **Strides Pharma Science Limited**
Strides House, Bilekahalli, Bannerghatta Road,
Bangalore - 560 076, Karnataka, Inde.

8510

8510

------ 100 x 190 mm ------

Back Page Printing

ARTWORK DETAIL LABEL

Product	**Melonax 7.5 & 15**				
Buyer/Country	SAFAL	Component	Pack Insert		
Dimension	**100 x 190 mm**			Pack	-------
New Item Code	**1040070**	Old Item Code	1039040		
Colour Shades	Pantone 506 U			No. of Colours	1

Change Control No.	PC-EM2/2019/130 - Record Number: 233784	Artwork Version	5.0
Design/Style	Front & Back Printing. To be supplied folded size of 50 x 23.75mm		
Substrate	60 GSM paper		
Special Instructions	PRINTING CLARITY TO BE CLEAR AND SHARP.		
Autocartonator Requirements	NA		

Caution to the printer: Before processing, please ensure that the ARTWORK received for printing is exactly in line with APPROVED ARTWORK provided to you. In case of any FONTS/DESIGN are Mis-matching with the APPROVED ARTWORK, please inform PDC for further action. **DO NOT MAKE ANY CHANGE TO THE ARTWORK WITHOUT WRITTEN INSTRUCTIONS FROM PDC.**