BLISS GVS PHARMA LTD.		
Product: DIAFORMINE 500/850 / 1000 mg TABLETS	Market: IMEX 2 (FR+PT+EN)	Size: (L x H) 456 x 260 mm
Component: Leaflet	Artwork code: 13006393-01	Pack Size:
Color Shades: Black	Date of Initiation:15-12-2018	Mfg Location: BLISS GVS
Barcode: N.A.	Substrate: 40 gsm Bible Paper	
Reason for Issue: Change in excipients		
Designer: Ramesh		Superceded Code: 13006393
G:\IMEX 3 Language Port-F-ENG\Diaformin/ Diaformin 500 C-L_B COMBINE LEAFLET		

18/11/2019

PATIENT INFORMATION LEAFLET: INFORMATION FOR THE USER

DIAFORMINE® 500 mg, coated tablets
DIAFORMINE® 850 mg, coated tablets
DIAFORMINE® 1000 mg, Sustained-release coated tablets

Metformin Hydrochloride

Please read this leaflet completely before taking this medicine as it contains important information:
- Keep this leaflet. You may need to read it again.
- Ask your doctor or pharmacist if you have any questions.
- This medication was prescribed for you only. Do not give it to others even if their symptoms are similar to yours as it may be harmful.
- Contact your doctor or pharmacist if you experience any side effects, including some not mentioned in this leaflet. See section 4.

What is in this leaflet?

1. What is DIAFORMINE®, coated tablets and when to use it?
2. What should I know before taking DIAFORMINE®, coated tablets?
3. How to take DIAFORMINE®, coated tablets?
4. What are the possible side effects?
5. How to store DIAFORMINE®, coated tablets?
6. Pack content and other information.

1. WHAT IS DIAFORMINE®, coated tablets AND WHEN TO USE IT?

Pharmacotherapeutic group—ATC code: A10BA02

DIAFORMINE® contains metformin a medication that belongs to the class of biguanidine drugs. It is used to treat diabetes.

Insulin, a hormone produced by the pancreas allows your body to recover the glucose (sugar) in the blood, which is then used to produce energy or stored for later use.

When you have diabetes, either your pancreas does not produce enough insulin or your body is not able to properly use the insulin produced. This results in a high glucose level in your blood. DIAFORMINE® helps lower your blood glucose level to as normal level as possible.

For overweight adults, long-term use of DIAFORMINE® can decrease the risk of complications associated to diabetes. DIAFORMINE® is associated with weight stability or modest weight loss.

DIAFORMINE® is used to treat type 2 diabetes (non-insulin-dependent diabetes) when diet and exercise alone were not enough to adequately control the blood glucose level. It is used more specifically in overweight patients.

Adults can be treated with DIAFORMINE® alone or with other diabetes medications (medication to be swallowed or insulin).

Children aged 10 or more and adolescents can be treated with DIAFORMINE® alone or with insulin.

2. WHAT SHOULD I KNOW BEFORE TAKING, coated tablets?

Never take DIAFORMINE®, coated tablets:
- If you are allergic (hypersensitive) to metformin or any of the other ingredients contained in this medication, see section 6.
- if you have liver problems.
- if you have severe deterioration of kidney function.
- if you have uncontrolled diabetes with, for example, severe hyperglycaemia (high blood glucose), nausea, vomiting, diarrhoea, rapid weight loss, lactic acidosis (see "Risk of lactic acidosis" below) or ketoacidosis. In case of ketoacidosis, substances called "ketone bodies" accumulate in the blood, which can lead to a diabetic pre-coma. Symptoms include stomach pain, fast and deep breathing, drowsiness, or unusually fruity-smelling breath.
- if you have lost too much water (dehydration) following persistent or severe diarrhoea, or vomiting several times in a row. Dehydration can lead to kidney problems that can put you at risk of lactic acidosis (see "Warnings and Precautions").
- if you have a severe infection, e.g., lungs, bronchi or kidney infection. Severe infections can lead to kidney problems that can put you at risk of lactic acidosis (see "Warnings and Precautions").
- if you take a treatment for acute heart failure or if you have recently had a heart attack. If you have severe blood circulation problems (such as shock) or if you have breathing difficulties. This can result in insufficient oxygen supply to the tissues putting you at risk of lactic acidosis (see "Warnings and Precautions").
- if you drink a lot of alcohol.

If any of the above applies to you, please consult your doctor before taking this medication.

Ask your doctor if:
- You need an X-Ray or CT scan that may require the injection of an iodine-based contrast product into the blood stream. •
- You are due to undergo major surgery.

Treatment with DIAFORMINE® should be interrupted a certain time before and after such examinations or surgeries. Your doctor will decide if a replacement treatment is required during that time. You must comply with the doctor's instructions.

Warnings and Precautions

Risk of lactic acidosis

DIAFORMINE® may cause a rare but serious side effect: lactic acidosis, especially if your kidneys do not function well. The risk of developing lactic acidosis increases with poorly controlled diabetes, serious infections, extended fasting or alcohol consumption, dehydration (see additional information below), liver problems and any other medical condition that leads to poor oxygen supply to parts of the body (e.g., severe acute heart disease).

If any of this applies to you, consult your doctor for further instructions.

Interrupt the DIAFORMINE® treatment for a short time if you have any medical condition that may be associated with dehydration (high loss of body fluids) such as severe vomiting, diarrhoea, fever, heat exposure, or if you have been drinking less fluids than usual. Talk to your doctor for further instructions.

Stop taking DIAFORMINE® and immediately contact a doctor or the nearest hospital if you experience lactic acidosis symptoms as this condition can lead to coma.

Lactic acidosis symptoms include:
- Vomiting.
- Stomach pains (abdominal pains).
- Muscle cramps.
- General feeling of being unwell with extreme fatigue.
- Difficulties breathing.
- Decreased body temperature and lower heart rate.

Lactic acidosis is a medical emergency that must be treated in hospital.

If you have to undergo major surgery, interrupt the DIAFORMINE® treatment at the time of the surgery and for a while following the procedure. Your doctor will decide when you should interrupt and resume the DIAFORMINE® treatment.

DIAFORMINE® on its own does not cause hypoglycaemia (low blood sugar level). However, there is a risk of hypoglycaemia if you combine DIAFORMINE® with other anti-diabetic medications (e.g., sulphonylurea, insulin, meglitinides). If you experience any symptoms associated with low-blood sugar, such as weakness, dizziness, increased sweating, fast heartbeat, blurred vision or trouble concentrating, drinking or eating something containing sugar should help you feel better.

During your treatment with DIAFORMINE®, your kidney function will be monitored at least once a year or more frequently if you are elderly and / or your kidney function has deteriorated.

Children

Non-applicable.

DIAFORMINE® coated tablets and other medications

If you are due to be injected an iodinated contrast agent for an X-Ray or CT scan you should stop taking DIAFORMINE® prior to or on the day of the injection. Your doctor will decide when you should stop and resume your treatment.

Inform your doctor if you take, have recently taken or are likely to take any other medication. You may need to have blood glucose or kidney function tests more frequently, or your doctor may have to adapt the DIAFORMINE® dosage. It is particularly important to report the medications listed below:
- medications that increase urine production (diuretics).
- medications used to treat pain and inflammation (NSAIDs or COX-2 inhibitors, such as ibuprofen and celecoxib).
- certain medications used to treat high blood pressure (ACE inhibitors and angiotensin II receptor antagonists).
- beta-2-adrenergic receptor agonists such as salbutamol or terbutaline (used to treat asthma).
- corticosteroids (used to treat various conditions, including severe skin inflammation or asthma).
- medications that may affect the amount of DIAFORMINE® in the blood, especially if you have reduced kidney function (such as verapamil, nifampicin, cimetidine, dolutegravir, ranolazine, trimethoprim, vandetanib, isavuconazonium, orizotinib and olaparib).
- other medications used to treat diabetes.

DIAFORMINE®, coated tablets with alcohol

Avoid consuming excessive amounts of alcohol when taking DIAFORMINE® as it may increase the risk of lactic acidosis (see "Warnings and Precautions").

Pregnancy and breastfeeding

During your pregnancy you may need insulin to treat your diabetes. Inform your doctor if you are pregnant, suspect or plan a pregnancy so your treatment may be adjusted.

This medication is not recommended if you are breastfeeding or plan to breastfeed.

Driving and using machines

DIAFORMINE® on its own does not cause hypoglycaemia (low blood sugar level). Therefore it should not affect your ability to drive or use machines.

However, be very careful if you take DIAFORMINE® with other anti-diabetic medications that can cause hypoglycaemia (e.g.: sulphonylurea, insulin, meglitinides). If you experience any symptoms associated with low-blood sugar, such as weakness, dizziness, increased sweating, fast heartbeat, blurred vision or trouble concentrating do not drive or use machines.

HOW TO TAKE DIAFORMINE®, coated tablets?

Always comply with the doctor's instructions. If in doubt consult your doctor or pharmacist.

DIAFORMINE® does not replace the benefits of a healthy lifestyle. Continue to follow your doctor's dietary advice and exercise regularly.

Recommended dosage

Children over 10 years old and adolescents: the starting dosage is usually 500 mg or 850 mg of DIAFORMINE® once a day. The maximum daily dose is 2000 mg in 2 or 3 separate doses.

Children between 10 and 12 should only be treated following specific advice from your doctor given the limited experience in this age group.

Adults: the starting dosage is usually 500 mg or 850 mg of DIAFORMINE® two or three times a day. The maximum daily dose is 3000 mg in 3 separate doses.

If you have impaired renal function, your doctor may prescribe a lower dosage.

If you also take insulin your doctor will tell you what dosage is better adapted.

Supervision
- Your blood sugar levels will have to be monitored regularly. Your doctor will adapt the DIAFORMINE® dosage based on the results of these tests. Be sure to consult your doctor regularly. This is especially important for children and adolescents or elderly patients.
- Your kidney function will have to be monitored at least once a year. Your doctor may require more frequent monitoring if you are an elderly patient or if your kidneys do not function normally.

How to take DIAFORMINE®, coated tablets

Take DIAFORMINE® during or after your meal to prevent any adverse effects on your digestion.

Do not crush or chew the tablets. Swallow each tablet with a glass of water.
- If you take 1 tablet per day, take it in the morning (at breakfast).
- If you take two separate doses per day, take one in the morning (at breakfast) and one in the evening (at dinner).
- If you take three separate doses per day, take one in the morning (at breakfast), one in the afternoon (at lunch) and one in the evening (at dinner).

After a while, if you have the impression that the effect of this medication is too strong or too weak, consult your doctor or pharmacist.

If you have taken more DIAFORMINE®, coated tablets than you should have:

Overdose of DIAFORMINE® can lead to the development of lactic acidosis. Lactic acidosis symptoms are non-specific, such as vomiting, stomach pains (abdominal pains), muscular cramps, general feeling of being unwell with extreme fatigue and difficulty breathing. Other symptoms include lower body temperature and slower heart rate. If you experience any of these symptoms consult a doctor immediately as lactic acidosis can lead to coma. Immediately stop taking DIAFORMINE® and contact the nearest hospital or doctor.

If you have forgotten to take DIAFORMINE®, coated tablets:

Do not take a double dose to compensate for the forgotten one. Take the next dose at the usual time.

If you continue taking DIAFORMINE®, coated tablets:

Non-applicable.

If you require further information pertaining to the use of this medication contact your doctor or pharmacist.

4. WHAT ARE THE POSSIBLE SIDE EFFECTS?

DIAFORMINE® may cause a very rare (can affect 1 in 10000 patients) but very serious side effect: lactic acidosis (see "Warnings and Precautions"). If you experience this, stop taking DIAFORMINE® and immediately contact the nearest hospital or doctor, as lactic acidosis can lead to coma.

As with all medications, some patients may experience side effects with this medication, although these are not systematic.

The following side effects may appear:

Most common side effects (may affect 1 in 10 patients):
- Digestive disorders, such as nausea, vomiting, diarrhoea, stomach ache (abdominal pain) and loss of appetite. These side effects are most often observed at the beginning of the treatment with DIAFORMINE®. Spreading the doses throughout the day and taking DIAFORMINE® during or just after the meal can help. If the symptoms persist, stop taking DIAFORMINE® and consult your doctor.

Common side effects (may affect 1 in 10 patients):
- Taste alteration.

Very rare side effects (may affect 1 in 10000 patients):
- Lactic acidosis: this is a very rare but serious complication, especially if your kidneys do not function well. Lactic acidosis symptoms are non-specific (see "Warnings and Precautions").
- Abnormal liver or hepatitis tests (liver inflammation that can cause fatigue, loss of appetite and weight with or without yellow colouring of the skin or white of the eyes). If this applies to you, stop taking DIAFORMINE® and consult your doctor.
- Skin reactions, such as redness of the skin (erythema), itching or itchy rash (hives).
- Decreased vitamin B12 in the blood.

Children and adolescents

Limited data in children and adolescents have shown that side effects were similar in nature and severity to those reported in adults.

Declaration of side effects

If you experience any side effects, even some not mentioned in this leaflet, inform your doctor or pharmacist. By declaring the side effects you help provide more information on the safety of this medication.

5. HOW TO STORE DIAFORMINE®, coated tablets?

Store below 30°C. Protect from light and moisture.

Keep out of reach and sight of children.

Do not use this medication after the expiry date on the box or blister indicated after "EXP". The expiry date refers to the last day of the month.

Do not dispose of in the sewer or with household wastes. Ask your pharmacist how to dispose of unused drugs. These measures allow protecting the environment.

6. PACK CONTENT AND OTHER INFORMATION
WHAT IS THE COMPOSITION OF DIAFORMINE®

DIAFORMINE® 500	DIAFORMINE® 850	DIAFORMINE® 1000
Each film coated tablet contains:	Each film coated tablet contains:	Each film coated sustained release tablet contains:
Metformin Hydrochloride 500 mg	Metformin Hydrochloride 850 mg	Metformin Hydrochloride 1000 mg
Excipients q.s.	Excipients q.s.	Excipients q.s

DIAFORMINE® 500: Microcrystalline Cellulose, Maize Starch, Povidone K30, Magnesium Stearate, Anhydrous colloidal silica (Aerosil), Isopropyl Alcohol, Dichloromethane, Wincoat WT-1001 White, Purified Talc.

DIAFORMINE® 850: Povidone (PvK30), Magnesium Stearate, Opadry White (03F58704).

DIAFORMINE® 1000: Hypromellose, Ethyl Cellulose, Magnesium Stearate, Anhydrous Colloidal Silica, Isopropyl Alcohol, Dichloromethane, Wincoat WT-1001 white, Purified Talc.

DIAFORMINE®, coated tablet, pack content and outer package

This medication is a white, circular & elongated, biconvex, film-coated tablet.

This medication comes in a box of 30 tablets.

This leaflet was last updated in 11/2019.

If you have any questions about this product or would like to report an adverse reaction contact us by phone : + 225 21 25 52 38

MA Holder & operator: **IMEX PHARMA**
Rose Belle Building, Unit 3, Rose Belle Business Park, Mauritius.

Manufacturer : **BLISS GVS PHARMA LTD.**
Dewan Udyog Nagar, Allyali, Palghar, Maharashtra - 401 404, INDIA.

10000300-01

NOTICE: INFORMATION DE L'UTILISATEUR

DIAFORMINE® 500 mg, comprimé pelliculé
DIAFORMINE® 850 mg, comprimé pelliculé
DIAFORMINE® 1000 mg, comprimé pelliculé à libération prolongée

Chlorhydrate de metformine

Veuillez lire attentivement cette notice avant de prendre ce médicament car elle contient des informations importantes pour vous.
- Gardez cette notice. Vous pourriez avoir besoin de la relire
- Si vous avez d'autres questions, interrogez votre médecin ou votre pharmacien
- Ce médicament vous a été personnellement prescrit. Ne le donnez pas à d'autres personnes. Il pourrait leur être nocif, même si les signes de leur maladie sont identiques aux vôtres
- Si vous ressentez un quelconque effet indésirable, parlez-en à votre médecin ou votre pharmacien. Ceci s'applique aussi à tout effet indésirable qui ne serait pas mentionné dans cette notice. Voir rubrique 4.

Que contient cette notice ?

1. Qu'est-ce que DIAFORMINE®, comprimé pelliculé et dans quels cas est-il utilisé ?
2. Quelles sont les informations à connaître avant de prendre DIAFORMINE®, comprimé pelliculé ?
3. Comment prendre DIAFORMINE®, comprimé pelliculé ?
4. Quels sont les effets indésirables éventuels ?
5. Comment conserver DIAFORMINE®, comprimé pelliculé ?
6. Contenu de l'emballage et autres informations.

1. QU'EST-CE QUE DIAFORMINE®, comprimé pelliculé ET DANS QUELS CAS EST-IL UTILISÉ ?

Classe pharmacothérapeutique – code ATC : A10BA02

DIAFORMINE® contient de la metformine. C'est un médicament utilisé pour traiter le diabète. Il appartient à la classe de médicaments appelés les biguanides.

L'insuline est une hormone produite par le pancréas et permettant à votre corps de récupérer le glucose (sucre) qui est dans le sang. Votre corps utilise le glucose pour produire de l'énergie ou le stocke pour l'utiliser plus tard.

Lorsque vous avez du diabète, votre pancréas ne produit pas suffisamment d'insuline ou votre corps n'est pas capable d'utiliser correctement l'insuline qu'il produit. Ceci aboutit à un taux de glucose élevé dans votre sang. DIAFORMINE® aide à faire baisser votre taux de glucose sanguin jusqu'à un niveau aussi normal qu'possible.

Si vous êtes un adulte en surpoids, la prise de DIAFORMINE® à long terme permet également d'aider à limiter les risques de complications associées au diabète. DIAFORMINE® est associé à une stabilité pondérale ou une modeste perte de poids.

DIAFORMINE® est utilisé pour traiter le diabète de type 2 (aussi appelé « diabète non insulino-dépendant ») lorsque le régime alimentaire et l'exercice physique seuls n'ont pas suffi à contrôler le taux de glucose sanguin. Il est particulièrement utilisé chez les patients en surpoids.

Les patients adultes pourront prendre DIAFORMINE® seul ou avec d'autres médicaments contre le diabète (des médicaments à avaler ou de l'insuline).

Les enfants de 10 ans et plus et les adolescents pourront prendre DIAFORMINE® seul ou avec l'insuline.

2. QUELLES SONT LES INFORMATIONS A CONNAITRE AVANT DE PRENDRE DIAFORMINE®, comprimé pelliculé ?

Ne prenez jamais DIAFORMINE®, comprimé pelliculé :
- Si vous êtes allergique (hypersensible) à la metformine ou à l'un des autres composants contenus dans ce médicament, mentionnés dans la rubrique 6
- Si vous avez des problèmes hépatiques
- Si vous souffrez d'une détérioration sévère de la fonction rénale
- Si vous avez un diabète non contrôlé avec, par exemple, une hyperglycémie sévère (taux élevé de glucose dans le sang), des nausées, des vomissements, une diarrhée, une perte de poids rapide, une acidose lactique (voir « Risque d'acidose lactique » ci-dessous) ou une acidocétose. En cas d'acidocétose, les substances appelées « corps cétoniques » s'accumulent dans le sang ce qui peut conduire à un pré-coma diabétique. Les symptômes comprennent des douleurs à l'estomac, une respiration rapide et profonde, une somnolence ou si votre haleine présente une odeur fruitée inhabituelle
- si vous avez perdu trop d'eau (déshydratation), par exemple suite à une diarrhée persistante ou sévère, ou si vous avez vomi plusieurs fois d'affilée. La déshydratation peut entraîner des problèmes de reins qui peuvent vous mettre en danger d'acidose lactique (voir « Avertissements et précautions »)
- si vous avez une infection grave, par exemple une infection des poumons ou des bronches, ou une infection des reins. Des infections sévères peuvent entraîner des problèmes de reins qui peuvent vous mettre en danger d'acidose lactique (voir « Avertissements et précautions »)
- si vous prenez un traitement contre l'insuffisance cardiaque aiguë ou si vous avez récemment eu une crise cardiaque, si vous souffrez de graves problèmes de circulation sanguine (tel un choc) ou si vous avez des difficultés respiratoires. Ceci peut entraîner un manque d'apport en oxygène des tissus, qui peut vous mettre en danger d'acidose lactique (voir « Avertissements et précautions »)
- si vous consommez beaucoup d'alcool

Si l'un des cas ci-dessus s'applique à vous, parlez-en à votre médecin avant de commencer à prendre ce médicament.

Veillez à demander conseil à votre médecin si :
- vous devez passer un examen radiographique ou un scanner qui pourrait nécessiter l'injection d'un produit de contraste à base d'iode dans votre circulation sanguine
- vous devez subir une intervention chirurgicale majeure

Vous devez arrêter de prendre DIAFORMINE® un certain temps avant et après de tels examens ou interventions chirurgicales. Votre médecin décidera si vous avez besoin ou non d'un traitement de remplacement pendant cette période. Il est important que vous suiviez précisément les instructions de votre médecin.

Avertissements et précautions

Risque d'acidose lactique

DIAFORMINE® peut provoquer un effet indésirable rare, mais très grave, appelé « acidose lactique », en particulier si vos reins ne fonctionnent pas correctement. Le risque de développer une acidose lactique augmente également en cas de diabète mal contrôlé, d'infections graves, de jeûne prolongé ou de consommation d'alcool, de déshydratation (voir informations complémentaires ci-dessous), de problèmes au foie et toutes autres affections médicales pour lesquelles une partie du corps reçoit un apport réduit en oxygène (les maladies cardiaques aiguës sévères, par exemple).

Si l'une de ces situations s'applique à vous, adressez-vous à votre médecin pour recevoir des instructions supplémentaires.

Arrêtez de prendre DIAFORMINE® pendant une courte période si vous souffrez d'une affection susceptible d'être associée à une déshydratation (perte importante de liquides corporels) tels des vomissements sévères, de la diarrhée, de la fièvre, une exposition à la chaleur ou si vous buvez moins de liquides que d'habitude. Adressez-vous à votre médecin pour obtenir des instructions supplémentaires.

Arrêtez de prendre DIAFORMINE® et contactez immédiatement un médecin ou l'hôpital le plus proche si vous présentez les symptômes d'une acidose lactique, car cette affection peut entraîner un coma.

Les symptômes de l'acidose lactique comprennent :
- Vomissements
- Maux d'estomac (douleurs abdominales)
- Crampes musculaires
- Sensation générale de malaise associée à une grande fatigue
- Difficultés à respirer
- Diminution de la température corporelle et du rythme cardiaque

L'acidose lactique est une urgence médicale qui doit être traitée à l'hôpital.

Si vous devez subir une opération chirurgicale importante, arrêtez de prendre DIAFORMINE® au moment de l'opération et pendant un certain temps après l'intervention. Votre médecin décidera du moment où vous devrez arrêter et reprendre votre traitement par DIAFORMINE®.

DIAFORMINE® à lui seul ne provoque pas d'hypoglycémie (taux de sucre dans le sang trop faible). Cependant, si vous prenez DIAFORMINE® en même temps que d'autres médicaments pour traiter le diabète qui peuvent engendrer une hypoglycémie (comme les sulfamides hypoglycémiants, l'insuline, les méglitinides), il y a un risque d'hypoglycémie. Si vous ressentez des symptômes d'hypoglycémie, tels que des tiraillements, des vertiges, une augmentation de la transpiration, des battements de cœur rapides, des troubles de la vision ou des difficultés à vous concentrer, boire ou manger quelque chose contenant du sucre devrait vous aider à vous sentir mieux.

Pendant votre traitement par DIAFORMINE®, votre médecin contrôlera votre fonction rénale au moins une fois par an ou plus fréquemment si vous êtes âgé(e) et/ou si votre fonction rénale s'est détériorée.

Enfants

Sans objet.

Autres médicaments et DIAFORMINE®, comprimé pelliculé

Si vous devez recevoir une injection d'un produit de contraste contenant de l'iode, par exemple, pour une radiographie ou un scanner, vous devrez arrêter de prendre DIAFORMINE® avant ou au moment de l'injection. Votre médecin décidera du moment où vous devrez arrêter et reprendre la prise de votre traitement par DIAFORMINE®

Informez votre médecin si vous prenez, avez récemment pris ou pourriez prendre tout autre médicament. Vous devrez peut-être passer plus fréquemment des tests de la glycémie ou de la fonction rénale ou votre médecin devra peut-être ajuster la posologie de DIAFORMINE®. Il est particulièrement important de signaler les médicaments suivants :
- les médicaments qui augmentent la production d'urine (diurétiques)
- des médicaments utilisés pour traiter la douleur et l'inflammation (AINS ou inhibiteurs de la COX-2, tels que l'ibuprofène et le célécoxib)
- certains médicaments utilisés pour traiter l'hypertension artérielle (inhibiteurs de l'ECA et antagonistes des récepteurs de l'angiotensine II)
- agonistes des récepteurs bêta-2-adrénergiques comme le salbutamol ou la terbutaline (utilisés pour traiter l'asthme)
- corticostéroïdes (utilisés pour traiter diverses affections, notamment des inflammations sévères de la peau ou l'asthme)
- les médicaments qui peuvent modifier la quantité de DIAFORMINE® dans le sang, en particulier si vous avez une fonction rénale réduite (comme le verapamil, la nifampicine, la cimétidine, le dolutégravir, la ranolazine, le trimethoprime, le vandetanib, l'isavuconazole, le crizotinib et l'olaparib)
- autres médicaments utilisés pour traiter le diabète

DIAFORMINE®, comprimé pelliculé avec de l'alcool

Éviter une consommation excessive d'alcool pendant la prise de DIAFORMINE®, car cela peut augmenter le risque d'acidose lactique (voir la rubrique « Avertissements et précautions »).

Front | English | English | French