INDOCO REMEDIES LTD.
Packaging Development Department
Packaging Material Specification

Item: Leaflet
A/W Code No.: COVBMLL01AESCI
Open Size: 110 X 255 mm
Folding Pattern: H-Unfolded: 255 mm Folded: 127.5 mm
W-Unfolded: 110 mm Folded: 27.5mm
GSM/Paper: 54 gsm maplitho paper
Colour: Black

Approved by:

PDD: _____ HOD: _____
Reason for Preparation: New Development for Baddi Plant I.

Pharma Code : 90 Std.

W:\Art-work\V\Viscid\Export\Ivory coast\Common\
New Prefex VBM\Viscid 200 ml Leaflet COVBMLL01AESCI.cdr
03.11.16 / 25.11.16/ 4.1.17

110 mm

255 mm

VISCID
Alumina, Magnesia and Simethicone Oral Suspension USP

PRESENTATION
Bottle of 200 ml

COMPOSITION
Each 5 ml contains:
Dried Aluminium Hydroxide Gel USP……400 mg
equivalent to Aluminium Hydroxide……...306 mg
Magnesium Hydroxide USP……100 mg
Simethicone Emulsion USP
equivalent to Simethicone USP……25 mg
Colour: Quinoline yellow WS

THERAPEUTIC CLASS
Drugs for acid related disorders; Antacids with anti-flatulents

PHARMACOLOGY
VISCID is a balanced mixture of two antacids and an anti-flatulent/antifoaming agent simethicone. The two antacids are magnesium hydroxide which is fast acting and aluminium hydroxide which is a slow acting antacid. The combination produces a fast onset of action and an increase in total buffering time. Aluminium hydroxide on its own is an astringent and may cause constipation. This effect is balanced by the effect of the magnesium hydroxide which is in common with other magnesium salts may cause diarrhoea.

PHARMACOKINETICS

None stated.

INDICATIONS
The symptomatic relief of:
1. Dyspepsia
2. Heartburn
3. Flatulence

CONTRAINDICATIONS
Should not be used in patients who are hypersensitive to any of the active substances or excipients, are severely debilitated or suffering from kidney failure, or hypophosphataemia.

WARNINGS AND PRECAUTIONS
In patients with renal impairment, plasma levels of both aluminium and magnesium increase. In these patients, a long-term exposure to high doses of aluminium and magnesium salts may lead to dementia, microcytic anemia.
Aluminum hydroxide may be unsafe in patients with porphyria undergoing hemodialysis.
This product contains sorbitol. Patients with rare hereditary problems of fructose intolerance should not take this medicine.

Pregnancy and Lactation
The safety of Viscid in pregnancy has not been established.
Magnesium is considered as compatible with lactation.

Pediatric Use
Children: As an appropriate proportion of the adult dose.

Effect on ability to drive and use machines
None stated.

ADVERSE EFFECTS
Gastrointestinal side-effects are uncommon.

DRUG INTERACTIONS
Viscid should not be taken simultaneously with other medicines as they may interfere with their absorption if taken within 1 hour.
Aluminium-containing antacids may prevent the proper absorption of cefdinir and cefpodoxime.
Aluminium hydroxide and citrates may result in increased aluminum levels, especially in patients with renal impairment.

DOSAGE AND ADMINISTRATION

For oral administration:

Adults: 5-10ml taken 20 minutes to 1 hour after meals and at bedtime or as required.

Children: As directed by the Physician.

Elderly: The normal adult dose is appropriate.

OVERDOSE
Serious symptoms are unlikely following overdosage.
Treatment of magnesium overdose: consider administration of IV Calcium Gluconate, rehydration and forced diuresis. In case of renal deficiency, haemodialysis or peritoneal dialysis is necessary.

STORAGE

Store below 30^{0}C.Protect from light

List of excipients
Carmellose Sodium (Sodium Carboxy Methyl Cellulose) BP , Sodium Methylhydroxybenzoate BP, Sodium Propylhydroxybenzoate BP, Propylene Glycol BP, Sorbitol Liquid (70%) (Non crystallizing) BP, Citric Acid Monohydrate BP, Saccharin Sodium BP, Colour Quinoline Yellow WS In- House , Fennel Flavour 50606 In-House, Bronopol (Bronidiol) BP, Purified Water BP

LIST I

Made in India by
INDOCO REMEDIES LTD.
Regd. Office: 166 C.S.T. Road,
Mumbai 400 098, INDIA

27.5 mm | 12 mm

110 mm

255 mm

VISCID
Suspension orale Aluminium, Magnésium et Simethicone USP

PRESENTATION
Bouteille de 200ml

COMPOSITION
Chaque 5 ml contient:
Séché d'hydroxyde d'aluminium Gel USP .....400 mg
Equivalent à de l'hydroxyde d'aluminium ..... 306 mg
L'hydroxyde de magnésium USP ....100 mg
Simethicone Emulsion USP
Equivalent à USP Simethicone ..... 25 mg
Couleur: jaune de quinoléine WS

CLASSE THERAPEUTIQUE
Troubles liés à l'hyperacidité; antiacides, anti-flatulents

PHARMACOLOGIE
VISCID est un mélange équilibré de deux antiacides et de Siméthicone,anti-flatulent/ Anti mousseux. Les deux antiacides sont l'hydroxyde de magnésium, qui est à action rapide et de l'hydroxyde d'aluminium qui est un anti-acide à action lente. La combinaison produit un début d'action rapide et une augmentation de la durée d'action au contact de la paroi digestive. L'hydroxyde d'aluminium seul e est un astringent et peut être source de constipation. Cet effet est compensé par celui de l'hydroxyde de magnésium qui associé à d'autres sels de magnésium peuvent provoquer des diarrhées.

PHARMACOCINETIQUES
Aucune donnée disponible**.**

INDICATIONS
VISCID est indiqué dans le soulagement symptomatique des troubles digestifs tels que:
1. Dyspepsie
2. Brûlures d'estomac
3. Flatulence

CONTRE-INDICATIONS
VISCID ne doit pas être utilisé chez les patients présentant une hypersensibilité à l'une des substances actives ou à un des excipients.
VISCID ne doit pas être utilisé chez les insuffisants rénaux, ou présentant une hypophosphatémie.

AVERTISSEMENTS ET PRÉCAUTIONS
Chez les patients souffrant d'insuffisance rénale, les concentrations plasmatiques d'aluminium et de magnésium augmentent. Chez ces patients, une exposition à long terme à de fortes doses de sels d'aluminium et de magnésium peut conduire à la démence, l'anémie microcytaire.
L'hydroxyde d'aluminium peut être dangereux chez les patients souffrants atteints de porphyrie et qui doivent **subir une hémodialyse**.
Ce produit contient du sorbitol. Les patients présentant des problèmes héréditaires rares d'intolérance au fructose ne doivent pas prendre ce médicament.

Grossesse et allaitement
La sécurité d'emploi de Viscid pendant la grossesse n'a pas été établie.
Le magnésium est compatible avec l'allaitement.

Utilisation en Pédiatrie
Enfants: Se conformer à la prescription du médecin

Conduite
Aucun effet sur la capacité de conduite de machines n'a été étudié

EFFETS INDÉSIRABLES
Les effets secondaires gastro-intestinaux sont rares

INTERACTIONS MEDICAMENTEUSES
Viscid ne doit pas être pris simultanément avec d'autres médicaments, car ils peuvent interférer l'absorption de Viscid s'il est pris en l'espace d' 1 heure.
Les antiacides contenant de l'aluminium peuvent empêcher la bonne absorption de la cefpodoxime et de la cefdinir.
L'hydroxyde d'aluminium et les citrates peuvent augmenter le taux d'aluminium, en particulier chez les patients présentant une insuffisance rénale.

POSOLOGIE ET MODE D'ADMINISTRATION

L'administration est orale:

Adultes: 5 à 10ml pris 20 minutes à 1 heure après les repas et au coucher ou selon les besoins.

Enfants: Selon prescription médicale.

Personnes âgées: Même dose que chez l'adulte.

SURDOSAGE
En cas de symptômes graves dus à un surdosage ;l'administration en IV de gluconate de calcium, la réhydratation et la diurèse forcée sont recommandées . En cas d'insuffisance rénale, l'hémodialyse ou la dialyse péritonéale est nécessaire.

CONSERVATION
Conserver en-dessous de 30°C. à l'abris de la lumière

Liste des excipients
Carmellose de sodium (Carboxyméthylcellulose) BP, Sodium Hydroxybenzoate de méthyle BP, propylhydroxybenzoate de Sodium BP, Propylène Glycol BP, sorbitol liquide (70%) (Non cristallisé) BP, monohydrate d'acide citrique BP, saccharine sodique BP, Couleur WS jaune de quinoléine, Saveur de Fenouil 50606, Bronopol (Bronidiol) BP, eau purifiée BP

Liste I

Fabriqué en Inde par

INDOCO REMEDIES LTD.
Siège social/Regd. Office:
166 C.S.T. Road, Mumbai 400 098
INDIA

COVBMLL01AESCI