


**84*43*103MM
TRAY PACK
30ML VIAL + 10ML WFI**

<p>Bactokil Imipenem & Cilastatin For Injection USP</p>	
<p>Single dose vial</p> <p style="text-align: center;">Bactokil Imipenem & Cilastatin For Injection USP (500 mg + 500 mg)</p> <p style="text-align: center;">For intravenous use only Pour usage intraveineuse</p> <p style="text-align: center; background-color: #e91e63; color: white; padding: 2px;">ÉCHANTILLON MÉDICAL MODÈLE-VENTE</p> <p style="text-align: right;">STERILE</p>	<p>Composition: Each vial contains : Imipenem (Sterile) USP eq. to Anhydrous Imipenem 500 mg Cilastatin Sodium (Sterile) USP eq. to Cilastatin 500 mg (Sterile Sodium Bicarbonate added as buffer) Dosage: As directed by the physician. Diluent supplied with this pack. M.L., B.No., Mfg. & Exp. Date of the diluent printed on the ampoule Storage: The dry powder should be stored below 30°C. Keep all medicines out of the reach of children. Direction for use: After Suspension vial contents must be transferred to 100 ml infusion solution. After reconstitution keep at room temperature 30°C for up to 4 hours or under refrigeration 5°C for up to 24 hours. Solution should not be frozen.</p> <p>Composition: Chaque flacon contient : Imipénème (stérile) USP éq. à Imipénème anhydre 500 mg Cilastatine sodique (stérile) USP éq. à Cilastatine 500 mg (Bicarbonate de sodium stérile ajouté comme tampon) Posologie: Selon les directives du médecin. Diluant fourni avec ce paquet. M.L., B.No., Mfg & Exp. Date du diluant imprimé sur l'ampoule Stockage: La poudre sèche doit être stocké en dessous de 30°C. Gardez tous les médicaments hors de la portée des enfants. Instructions: Après la suspension, le contenu du flacon doit être transféré dans 100 ml de solution pour perfusion. Après reconstitution garder dans la chambre température 30°C jusqu'à 4 heures ou sous réfrigération à 5°C jusqu'à 24 heures. La solution ne doit pas être gelée.</p>
<p>Bactokil Imipénème et cilastatine pour injection USP (500 mg + 500 mg)</p> <p style="text-align: center;">For intravenous use only Pour usage intraveineux</p> <p style="text-align: center; background-color: #e91e63; color: white; padding: 2px;">ÉCHANTILLON MÉDICAL MODÈLE-VENTE</p> <p style="text-align: right;">STÉRILE</p>	
<p>Warning: If any foreign particle is visible in the bottle after reconstitution, please do not use the solution.</p> <p>Attention: si une particule étrangère est visible dans la bouteille après reconstitution, veuillez ne pas utiliser la solution.</p>  <p style="text-align: center; background-color: #e91e63; color: white; padding: 2px;">DO NOT EXCEED THE PRESCRIBED DOSE NE PAS DÉPASSER LA DOSE PRESCRITE</p> <p>Mfg. Lic. No. : HP/DRUGS/08/35-36 Batch No. : Mfg. Date : Exp. Date :</p> <p>Manufactured in India / Fabriqué en Inde .</p>  <p>MYRA HEALTHCARE Office No.1, Shree OM Sadguru CHS, Plot No. 122, Sector 50-E, Seawood, Nerul (W), Navi Mumbai (India)</p>	

CHECK LIST

Brand Name <input type="text"/>	Generic Name <input type="text"/>	I.P./B.P./U.S.P. <input type="text"/> <small>(Generic Name & Composition)</small>	Flap Brand <input type="text"/>	Mfg. Address <input type="text"/>	Foil Size <input type="text"/>	Print Area <input type="text"/>
Marketed by <input type="text"/>	Mfg. Lic. No. <input type="text"/>	Rx/NRx <input type="text"/> <small>Red Line</small>	Carton Size <input type="text"/>	Packing Style <input type="text"/>	Print Repeat <input type="text"/>	Total Print Area <input type="text"/>
Composition <input type="text"/> <small>(as per FDA)</small>	Mode of Adm. <input type="text"/> <small>(I.M./I.V/S.C.)</small>	Schedule <input type="text"/> <small>(G/H/H1)</small>	Supply / MRP <input type="text"/>	Warnings <input type="text"/>		

Signature (QA)
